

Gender differences in coronary functional abnormalities

Kensuke Nishimiya, Hiroaki Shimokawa

Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan

Received 2 September 2020; accepted 17 September 2020

Summary. The ISCHEMIA trial demonstrated that coronary revascularization strategies may not necessarily improve the long-term prognosis of patients with obstructive coronary artery disease, suggesting the significance of the remaining coronary functional abnormalities. Recently, more attention has been paid to ischemia with no obstructive coronary artery (INOCA) disease, due to the poor prognosis as showed in several cohorts. Coronary microvascular dysfunction (CMD) and epicardial coronary spasm have been proposed as the main causes of INOCA. While gender difference was not significant in the prognosis of patients with epicardial spasm, a poor prognosis was identified in women with CMD. In the last decade, efforts have been made to clarify the connection between these coronary functional abnormalities and an impaired prognosis. In this article, we will briefly review the recent advances in gender medicine regarding coronary functional abnormalities, including CMD and epicardial spasm.

Key words. Coronary microvascular dysfunction, microvascular spasm, coronary artery spasm, vasospastic angina.

Differenze di genere nelle anomalie funzionali coronariche

Riassunto. Lo studio ISCHEMIA ha dimostrato che le strategie di rivascularizzazione coronarica non necessariamente migliorano la prognosi a lungo termine dei pazienti con malattia coronarica ostruttiva, suggerendo l'importanza delle anomalie funzionali coronariche. Recentemente è stata prestata maggiore attenzione all'ischemia con malattia coronarica non ostruttiva (INOCA) a causa della prognosi sfavorevole che ha dimostrato in diverse coorti di pazienti. La disfunzione microvascolare coronarica (CMD) e lo spasmo coronarico epicardico sono stati proposti come le principali cause di INOCA. Sebbene la differenza di genere non fosse significativa nella prognosi dei pazienti con spasmo epicardico, nelle donne con CMD è stata identificata una prognosi sfavorevole. Nell'ultimo decennio sono stati compiuti sforzi per chiarire la connessione tra le anomalie funzionali coronariche e una prognosi compromessa. In questo articolo esamineremo brevemente i recenti progressi nella medicina di genere per quanto riguarda le anomalie funzionali coronariche, tra cui CMD e spasmo epicardico.

Parole chiave. Disfunzione microvascolare coronarica, spasmo microvascolare, spasmo dell'arteria coronaria, angina vasospastica.

Introduction

Recently, the Objective Randomized Blinded Investigation with optimal medical Therapy of Angioplasty in Stable Angina (ORBITA)¹ and the International Study of Comparative Health Effectiveness with Medical and Invasive Approaches (ISCHEMIA)² trials demonstrated that coronary revascularization strategies may not necessarily improve the clinical manifestations or the long-term prognosis of patients with stable obstructive coronary artery disease (CAD), thus suggesting the significance of the remaining coronary functional abnormalities. Another landmark study, the WISE (Women's Ischemia Syndrome Evaluation) trial, had previously reported that approximately 50-70% female patients with suspected angina had an angiographically insignificant coronary stenosis.³ This disorder, called ischemia with no obstructive coronary artery (INOCA), should be given much attention, because it has been emerging for the poor prognosis demonstrated in several cohorts.⁴⁻⁶ The WISE study estimated that at least 3-4 million patients in the USA are diagnosed with INOCA.^{3,4,7,8} Although multiple mechanisms may be involved in the pathogenesis of INOCA, coronary microvascular dysfunction (CMD) and epicardial coronary spasm are considered to be the leading causes of the disorder.⁹⁻¹³ In particular, CMD is more prevalent in women than in men, with poor clinical outcomes.¹³ In this article, we will briefly review the recent reports on the gender differences in coronary functional abnormalities (Figure).

Gender differences in CMD

Microvascular angina (MVA) is the term that is widely used to describe patients with symptomatic CMD.¹⁴ The consensus document by the Coronary Vasomotor Disorders International Study Group (COVADIS) proposed that MVA is diagnosed based on the following criteria:¹⁴ (1) presence of symptoms suggestive of myocardial ischemia; (2) objective documentation of myocardial ischemia by currently available techniques; (3) absence of obstructive CAD (<50% coronary diameter reduction and/or fractional flow reserve (FFR) >0.80), and (4) con-

firmation of a reduced coronary blood flow reserve and/or inducible microvascular spasm (MVS). CMD variably comprises several major pathophysiological mechanisms, including enhanced vascular smooth muscle contraction (MVS), impaired microvascular dilatation, and vascular remodeling associated with structural heart disease.^{10,15} MVS is diagnosed through the reproduction of angina symptoms accompanied by ischemic ECG changes, but with no epicardial spasm during spasm provocation testing.¹³ Additionally, interventional diagnostic approaches with coronary flow reserve (CFR, ≤ 2.0 or 2.5),¹⁶ index of microcirculatory resistance (IMR, >25),¹⁷ or thrombolysis in myocardial infarction frame count (TIMI frame count, >25)¹⁸ are also useful for making a diagnosis of CMD.

Gender differences in the epidemiology and prognosis of CMD


It has been consistently demonstrated that CMD is more prevalent in females than in males, especially in postmenopausal women (Figure).¹⁹⁻²¹ Indeed, in the Coronary Microvascular Angina (CorMicA) study, female sex was more correlated with CMD versus male [odds ratio, 2.7 (95% CI: 0.9-7.9)].²² In the WISE study, with a median of 5.4 follow-up years, CMD (determined by CFR <2.3) was closely associated with Major Adverse Cardiac Events (MACE), including hospitalization for acute coronary syndrome, heart failure, stroke, or cardiovascular death, with 53% of sudden cardiac death.³ Physiological levels of estradiol (E2) exert vasculoprotective and anti-inflam-

matory effects.²³ It has been speculated that women are closely associated with inflammatory markers [eg., high-sensitive c-reactive protein (hs-CRP)], and that inflammatory diseases occur more frequently in menopausal women than in men.²⁴ Sex differences in vasoreactivity to mental stress should also be mentioned. Peripheral vasoconstriction induced by a public speaking stressor is associated with stress-induced myocardial (microvascular) ischemia in women, but not in men.²⁵ Importantly, a positron emission tomography (PET) study showed that both sexes were at high risk for MACE (Figure).²⁶

In general, patients with MVA exhibit better angina summary score than those with vasospastic angina (VSA).²² Regarding the gender variation in the clinical presentation of CMD, young and middle-aged women are more likely to present with 'atypical' angina symptoms.²⁷ These atypical symptoms of MVA could occur at low heart rates, due to mental arousal or palpitation.¹⁰ Although known atherosclerotic risk factors, such as age, diabetes mellitus, hypertension, and dyslipidemia, have been identified as possible causes of CMD,^{11,12} these risk factors are not always predictive.²⁸ Thus, this point still remains to be addressed in future studies.

Heterogeneity of gender differences in the various diagnostic markers of CMD

Regarding the diagnosis of MVS during acetylcholine (ACh) provocation testing,¹⁴ Ong et al. found that the prevalence of female patients was higher in MVS than



Gender differences in coronary functional abnormalities are summarized. Female patients are more likely to have CMD and diffuse epicardial spasm. The prognostic effects of gender differences and the gender-specific treatment for coronary functional abnormalities remain to be elucidated. Numbers represent the corresponding reference numbers.

in VSA [169/205 (82%) in MVS vs 155/283 (55%) in VSA].²⁰ They also demonstrated that female sex was an independent predictor for MVS (OR = 4.2, 95% CI: 3.1-5.5, $p < 0.001$) (Figure).²⁹ In order to make a more precise diagnosis of MVS, we proposed to use the myocardial lactate extraction ratio, which can be calculated as the ratio of coronary arterial vs venous difference in lactate concentration to the arterial concentration during spasm provocative testing.^{9,30-32} In our study, female prevalence was comparable between non-MVS and MVS groups [50/132 (38%) vs 32/66 (49%) patients].³² Ethnicity may also be involved in the gender difference of the prevalence of MVS.

Physiological indexes, such as low CFR and high IMR, have been used as gold standards for the diagnosis of CMD (Figure).¹⁰⁻¹⁴ These indexes are measured in the left anterior descending coronary artery via inserted pressure/temperature sensor-mounted guide wire. Briefly, CFR is calculated using thermodilution as resting mean transit time divided by hyperemic mean transit time, whereas IMR is the product of distal coronary pressure at maximal hyperemia multiplied by the hyperemic mean transit time.³³ Ford et al. clearly demonstrated that female prevalence was higher in the MVA group based on the criteria of either CFR < 2 , IMR ≥ 25 or MVS for MVA [women, 61/78 (78%)].²² Kobayashi et al. reported that no gender difference was noted in IMR, whereas CFR was lower in women than in men (Figure).³⁴ Lee et al. reported that no remarkable gender difference was noted among 230 patients with intermediate coronary stenosis when sorting them into the quadrant according to CFR (a cut-off value of 2) and IMR (a cut-off value of 23).³⁵ We also demonstrated that, among INOCA patients, female sex was not a significant contributing factor to the presence of high IMR ≥ 18 , a prognostic marker for INOCA (OR = 0.51, 95% CI: 0.3-1.0 in the multivariable COX analysis).³⁶ A recent study by Chung et al. showed that, in patients deferred by FFR, CFR related to mean transit time was lower in women than in men, with no differences in IMR.³⁷ However, in this study, long-term clinical outcomes during a 5-year follow-up were better in women than in men (Figure).³⁷

PET allows the CFR quantification of myocardial blood flow (MBF, ml/min/g) of tissue both at rest and during maximal coronary vasodilatation.³⁸ Murthy et al. demonstrated that, among patients with no visual evidence of CAD, CMD determined by CFR < 2 on PET existed equally between men and women (51% and 54%), and showed no gender difference in the long-term prognosis.²⁶ Gupta et al. reported an interesting finding, namely that women were the majority of the discordant group with impaired CFR, but with preserved maximum MBF [608/873 (70%) patients].³⁹ They considered that this discordant group may be a target for the initiation of lifestyle or pharmacological preventive therapies, and

may aid the gender differences in the cardiovascular outcome.³⁹ Cardiac magnetic resonance (CMR) imaging could be an option for making the diagnosis of CMD in women, since it can prevent radiation exposure.⁴⁰

Gender differences in the treatment of CMD

Efforts to establish the optimal medical therapy for CMD have just begun, and several pharmacological strategies are under investigations. The currently proposed anti-anginal agents for CMD include β -blockers, calcium-channel blockers (CCB), angiotensin-converting enzyme inhibitors (ACEi)/angiotensin receptor blockers, nicorandil, ranolazine, ivabradine, and trimetazidine.¹¹ Among them, ACEi have been reported to improve CFR and angina frequency in female CMD patients.¹¹ Despite the high prevalence of menopausal women in CMD,⁴¹ the estrogen replacement therapy is not fully recommended for CMD, because it is possibly associated with increased risks of cardiovascular disease and breast cancer.⁴²

Female patients with INOCA have been reported to have impaired exercise capability (Figure). For instance, it was reported that peak VO_2 was lower in patients with CMD than in controls (17.3 vs 27.3 ml/kg/min).⁴³ Exercise and rehabilitation could improve the clinical manifestations in CMD patients. The COVADIS group started the international prospective cohort study for CMD patients.⁴⁴ This ongoing study is expected to provide novel robust evidence on gender differences in the ethnicity, clinical manifestations, and management of MVA patients in the near future.

Gender differences in vasospastic angina (VSA)

It has been believed for a long time that there are substantial ethnic differences in the prevalence of vasospastic angina (VSA) due to epicardial coronary artery spasm. Indeed, patients with acute myocardial infarction who were positive for spasm provocative testing were more common in Japan than among Caucasians (80% vs 37%).⁴⁵ However, recent studies demonstrated that such ethnic difference in the prevalence of VSA is much smaller than anticipated.^{46,47} In addition to CMD, this coronary functional abnormality by epicardial coronary spasm is important in order to better understand the pathogenesis of INOCA.

Gender differences in epidemiology and prognosis of VSA

A multicentre study conducted in Japan showed that 921 out of 2,251 angina patients were diagnosed with VSA.⁴⁸ In this study, patient distribution peaked at the age of 60-69, but was commonly documented in the under-60s.⁴⁸ Similar trends have been reported in the European

studies.^{29,49} We also have recently addressed this issue of ethnic difference in VSA.⁴⁷ Although female patients with VSA were not prevalent in both Japanese and Caucasians populations (13-22 %), the frequency of women was significantly higher among Caucasians than Japanese (Figure).⁴⁷ The Abnormal COronary VAsomotion (ACOVA) study reported that the epicardial spasm was more frequent in women than in men (77% vs 23%).⁴⁹ Similarly, Ford et al. reported that female patients with VSA accounted for 60%, while female sex was not a significant associate of VSA in multivariable analysis.²² However, in these studies, most of the participants were female, thus caution should be used when considering gender differences in VSA in the real world setting. Although the involvement of genetic factors in VSA is still being debated, Ong et al. reported that, among women with INOCA, a positive family history was more predominant in women than in men (61% vs 45%, $p = 0.0001$).⁵⁰ Furthermore, anginal symptoms of premenopausal female patients with VSA were closely associated with endogenous estrogen levels, where the frequency of angina attack peaked between late luteal and menstrual phases.⁵¹

Through the multicenter registry of the Japanese Coronary Spasm Association (JCSA), we demonstrated that no significant gender difference was noted in the 5-year MACE-free survival (Figure).⁵² We also reported that the hazard ratio of male sex was 1.1 (CI: 0.6-1.8, $p = 0.79$).⁵³ Furthermore, among female patients with VSA, the long-term prognosis was worst in the young female arm with a smoking behavior.⁵² These results indicate the importance of a gender-specific management of VSA. We also demonstrated that a Rho-kinase activity >1.2 could be a prognostic maker for VSA, for which a higher male prevalence was noted among patients with enhanced Rho-kinase activity.^{54,55} Most recently, we revealed that CMD co-existed especially in female VSA patients, a situation in which Rho-kinase activation may substantially be involved (Figure).⁵⁶

Gender differences in the spasm provocation testing of VSA

The Japanese Circulation Society (JCS) guidelines and the latest European consensus document highly recommend spasm provocation testing with ACh when a subject is negative for non-invasive VSA evaluation, but is still suspected to have coronary spasm based on symptoms.^{13,48} Ergonovine provocation testing is another option, when ACh is contraindicated due to comorbid bronchial asthma or severe atrioventricular conduction disorder.⁵⁷ Sueda et al. reported that female patients were unlikely to be sensitive to ergonovine (provocation rate; 96.7% with ACh vs 32.8% with ergonovine) (Figure).⁵⁸ Furthermore, Sato et al. reported that there was a distinct sex difference in the prevalence of diffuse/focal spasm;

60/33% in women vs 40/67% in male, respectively (Figure).⁵⁹ In the JCSA registry study, female sex was an independent factor for provocation testing related to ventricular arrhythmias.⁶⁰ Anatomical anomalies, especially myocardial bridging (MB),⁶¹ are the preferred site of epicardial coronary spasm. Although anatomical MB is more common in men than in women, there was no sex difference in functional abnormalities and flow disturbance due to MB.⁶²

Gender differences in the treatment of VSA

CCBs are the first-line therapy for VSA, since these drugs have been shown to markedly reduce angina symptom⁶³⁻⁶⁵ and suppress coronary spasm.^{63,64} CCBs have been an independent determinant for preventing MACE in VSA.^{63,64} The prognostic effects of nitrates on VSA remain unclear.⁶⁶ Although it has been considered that an insufficient medical therapy could result in a poorer prognosis in female patients, the recent JCSA registry study confirmed that the prescription of CCBs was adequately done in female VSA patients.⁵² The one-year follow-up result of the CorMicA study showed that the gender difference was unclear in the effectiveness of an optimized management for INOCA, based on the interventional diagnostic approaches.⁶⁷ We previously demonstrated that estrogen down-regulates the Rho-kinase activity,⁶⁸ a central molecular mechanism of coronary spasm.^{9,69} The JCS guidelines state that the estrogen replacement therapy is recommended in postmenopausal women as Class 2b.⁴⁸ Coronary plaque erosion has recently attracted more attention than ever before, because this phenomenon is the second leading cause of acute coronary syndrome (ACS),⁷⁰ for which epicardial coronary spasm could be a trigger.⁷¹ In an autopsy study, the prevalence of coronary erosion was higher than that of plaque rupture in female.⁷² Further studies are needed to address the underlying mechanism of this major subset of ACS.

Key messages

- Coronary microvascular dysfunction (CMD) and epicardial coronary spasm have been proposed as the main causes of INOCA.
- The prevalence of female patients is prominent in ischemia with no obstructive coronary artery (INOCA), particularly in CMD.
- There are plausible connections between gender differences and disease manifestations between CMD and epicardial coronary spasm.
- It is important to address gender differences in these coronary functional abnormalities in INOCA patients.

Conclusions

The prevalence of female patients is prominent in INOCA, particularly in CMD. There are plausible connections between gender differences and disease manifestations between CMD and epicardial coronary spasm. While seeking the perfect explanation of the mechanism and the optimization of the management of INOCA, it is important to address gender differences in these coronary functional abnormalities in INOCA patients. An international standardization strategy and expert consensus-based evidence are required for the promotion of women's cardiac health in the future.

References

- Al-Lamee R, Thompson D, Dehbi HM, Sen S, Tang K, Davies J, et al. Percutaneous coronary intervention in stable angina (ORBITA): a double-blind, randomized controlled trial. *Lancet*. 2018; 391(10115):31-40.
- Maron DJ, Hochman JS, Reynolds HR, Bangalore S, O'Brien SM, Boden WE, et al. Initial invasive or conservative strategy for stable coronary disease. *N Engl J Med*. 2020;382(15):1395-407.
- Pepine CJ, Anderson RD, Sharaf BL, Reis SE, Smith KM, Handberg EM, et al. Coronary microvascular reactivity to adenosine predicts adverse outcome in women evaluated for suspected ischemia results from the National Heart, Lung and Blood Institute WISE (Women's Ischemia Syndrome Evaluation) study. *J Am Coll Cardiol*. 2010;55(25):2825-32.
- Johnson BD, Shaw LJ, Pepine CJ, Reis SE, Kelsey SE, Sopko G, et al. Persistent chest pain predicts cardiovascular events in women without obstructive coronary artery disease: results from the NIH-NHLBI-sponsored Women's Ischaemia Syndrome Evaluation (WISE) study. *Eur Heart J*. 2006;27(12):1408-15.
- Jespersen L, Hvelplund A, Abildstrøm SZ, Pedersen F, Galatius S, Madsen JK, et al. Stable angina pectoris with no obstructive coronary artery disease is associated with increased risks of major adverse cardiovascular events. *Eur Heart J*. 2012;33(6):734-44.
- Maddox TM, Stanislawski MA, Grunwald GK, Bradley SM, Ho PM, Tsai TT, et al. Non-obstructive coronary artery disease and risk of myocardial infarction. *JAMA*. 2014;312(17):1754-63.
- von Mering GO, Arant CB, Wessel TR, McGorray SP, Bairey Merz CN, Sharaf BL, et al. Abnormal coronary vasomotion as a prognostic indicator of cardiovascular events in women: results from the National Heart, Lung, and Blood Institute-Sponsored Women's Ischemia Syndrome Evaluation (WISE). *Circulation*. 2004;109(6):722-5.
- Bairey Merz CN, Shaw LJ, Reis SE, Bittner V, Kelsey SE, Olson M, et al. Insights from the NHLBI-Sponsored Women's Ischemia Syndrome Evaluation (WISE) Study: Part II: gender differences in presentation, diagnosis, and outcome with regard to gender-based pathophysiology of atherosclerosis and macrovascular and microvascular coronary disease. *J Am Coll Cardiol*. 2006;47(3 Suppl):S21-9.
- Shimokawa H. 2014 Williams Harvey Lecture: importance of coronary vasomotion abnormalities—from bench to bedside. *Eur Heart J*. 2014;35(45):3180-93.
- Crea F, Camici PG, Bairey Merz CN. Coronary microvascular dysfunction: an update. *Eur Heart J*. 2014;35(17):1101-11.
- Bairey Merz CN, Pepine CJ, Walsh MN, Fleg JL. Ischemia and no obstructive Coronary Artery Disease (INOCA): developing evidence-based therapies and research agenda for the next decade. *Circulation*. 2017;135(11):1075-92.
- Ford TJ, Rocchiccioli P, Good R, McEntegart M, Eteiba H, Watkins S, et al. Systemic microvascular dysfunction in microvascular and vasospastic angina. *Eur Heart J*. 2018;39(46):4086-97.
- Kunadian V, Chieffo A, Camici PG, Berry C, Escaned J, Maas AHAM, et al. An EAPCI expert consensus document on ischaemia with non-obstructive coronary arteries in collaboration with European society of cardiology working group on coronary pathophysiology & microcirculation endorsed by coronary vasomotor disorders international study group. *Eur Heart J* 2020. doi: <https://doi.org/10.1093/eurheartj/ehaa503>.
- Ong P, Camici PG, Beltrame JF, Crea F, Shimokawa H, Sechtem U, et al. International standardization of diagnostic criteria for microvascular angina. *Int J Cardiol*. 2018;250:16-20.
- Camici PG, d'Amati G, Rimoldi O. Coronary microvascular dysfunction: mechanisms and functional assessment. *Nat Rev Cardiol*. 2015;12(1):48-62.
- Reis SE, Holubkov R, Lee JS, Sharaf B, Reichek N, Rogers WJ, et al. Coronary flow velocity response to adenosine characterizes coronary microvascular function in women with chest pain and no obstructive coronary disease. Results from the pilot phase of the Women's Ischemia Syndrome Evaluation (WISE) study. *J Am Coll Cardiol*. 1999; 33(6):1469-75.
- Fearon WF, Kobayashi Y. Invasive assessment of the coronary microvasculature: the index of microcirculatory resistance. *Circ Cardiovasc Interv*. 2017;10(12):e005361.
- Gibson CM, Cannon CP, Daley WL, Dodge JT Jr, Alexander B Jr, Marble SJ, et al. TIMI frame count: a quantitative method of assessing coronary artery flow. *Circulation*. 1996;93(5):879-88.
- Masumoto A, Mohri M, Takeshita A. Three-year follow-up of the Japanese patients with microvascular angina attributable to coronary microvascular spasm. *Int J Cardiol*. 2001;81(2-3):151-6.
- Ong P, Athanasiadis A, Borgulya G, Vokshi I, Bastiaenen R, Kubik S, et al. Clinical usefulness, angiographic characteristics, and safety evaluation of intracoronary acetylcholine provocation testing among 921 consecutive white patients with unobstructed coronary arteries. *Circulation*. 2014;129(17):1723-30.
- Murakami H, Urabe K, Nishimura M. Inappropriate microvascular constriction produced transient ST-segment elevation in patients with syndrome X. *J Am Coll Cardiol*. 1998;32(5):1287-94.
- Ford TJ, Yii E, Sidik N, Good R, Rocchiccioli P, McEntegart M, et al. Ischemia and no obstructive coronary artery dis-

- ease: prevalence and correlates of coronary vasomotion disorders. *Circ Cardiovasc Interv.* 2019;12(12):e008126.
23. Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. *N Engl J Med.* 1999;340(23):1801-11.
 24. Mommersteeg PMC, Naudé PJW, Bagijn W, Widdershoven J, Westerhuis BWJMM, Schoemaker RG. Gender differences in associations of depressive symptoms and anxiety with inflammatory markers in patients with non-obstructive coronary artery disease. *J Psychosom Res.* 2019;125:109779.
 25. Al-Badri A, Kim JH, Liu C, Mehta PK, Quyyumi AA. Peripheral microvascular function reflects coronary vascular function. *Arterioscler Thromb Vasc Biol.* 2019;39(7):1492-500.
 26. Murthy VL, Naya M, Taqueti VR, Foster CR, Gaber M, Hainer J, et al. Effects of sex on coronary microvascular dysfunction and cardiac outcomes. *Circulation.* 2014;129(24):2518-27.
 27. Biddle C, Fallavollita JA, Homish GG, Giovino GA, Orom H. Gender differences in symptom misattribution for coronary heart disease symptoms and intentions to seek health care. *Women Health.* 2020;60(4):367-81.
 28. Waheed N, Elias-Smale S, Malas W, Maas AH, Sedlak TL, Tremmel J, et al. Sex differences in non-obstructive coronary artery disease. *Cardiovasc Res.* 2020;116(4):829-40.
 29. Aziz A, Hansen HS, Sechtem U, Prescott E, Ong P. Sex-related differences in vasomotor function in patients with angina and unobstructed coronary arteries. *J Am Coll Cardiol.* 2017;70(19):2349-58.
 30. Sun H, Mohri M, Shimokawa H, Usui M, Urakami L, Takeshita A. Coronary microvascular spasm causes myocardial ischemia in patients with vasospastic angina. *J Am Coll Cardiol.* 2002;39(5):847-51.
 31. Mohri M, Shimokawa H, Hirakawa Y, Masumoto A, Takeshita A. Rho-kinase inhibition with intracoronary fasudil prevents myocardial ischemia in patients with coronary microvascular spasm. *J Am Coll Cardiol.* 2003;41(1):15-9.
 32. Odaka Y, Takahashi J, Tsuburaya R, Nishimiya K, Hao K, Matsumoto Y, et al. Plasma concentration of serotonin is a novel biomarker for coronary microvascular dysfunction in patients with suspected angina and unobstructive coronary arteries. *Eur Heart J.* 2017;38(7):489-96.
 33. Ford TJ, Ong P, Sechtem U, Beltrame J, Camici PG, Crea F, et al. Assessment of vascular dysfunction in patients without obstructive coronary artery disease: why, how, and when. *J Am Coll Cardiol Interv.* 2020;13(16):1847-64.
 34. Kobayashi Y, Fearon WF, Honda Y, Tanaka S, Pargaonkar V, Fitzgerald PJ, et al. Effect of sex differences on invasive measures of coronary microvascular dysfunction in patients with angina in the absence of obstructive coronary artery disease. *J Am Coll Cardiol Interv.* 2015;8(11):1433-41.
 35. Lee JM, Jung JH, Hwang D, Park J, Fan Y, Na SH, et al. Coronary flow reserve and microcirculatory resistance in patients with intermediate coronary stenosis. *J Am Coll Cardiol.* 2016;67(10):1158-69.
 36. Suda A, Takahashi J, Hao K, Kikuchi Y, Shindo T, Ikeda S, et al. Coronary functional abnormalities in patients with angina and nonobstructive coronary artery disease. *J Am Coll Cardiol.* 2019;74(19):2350-60.
 37. Chung JH, Lee KE, Lee JM, Her AY, Kim CH, Choi KH, et al. Effect of sex difference of coronary microvascular dysfunction on long-term outcomes in deferred lesions. *J Am Coll Cardiol Interv.* 2020;13(14):1669-79.
 38. Camici PG, Rimoldi OE. The clinical value of myocardial blood flow measurement. *J Nucl Med.* 2009;50(7):1076-87.
 39. Gupta A, Taqueti VR, van de Hoef TP, Bajaj NS, Bravo PE, Murthy VL, et al. Integrated noninvasive physiological assessment of coronary circulatory function and impact on cardiovascular mortality in patients with stable coronary artery disease. *Circulation.* 2017;136(24):2325-36.
 40. Wei J, Bakir M, Darounian N, Li Q, Landes S, Mehta PK, et al. Myocardial scar is prevalent and associated with subclinical myocardial dysfunction in women with suspected ischemia but no obstructive coronary artery disease: from the women's ischemia syndrome evaluation-coronary vascular dysfunction study. *Circulation.* 2018;137(8):874-6.
 41. Kaski JC. Cardiac syndrome X in women: the role of oestrogen deficiency. *Heart.* 2006;92 Suppl 3(Suppl 3):iii5-9.
 42. Mosca L, Benjamin EJ, Berra K, Bezanson JL, Dolor RJ, Lloyd-Jones DM, et al. Effectiveness-based guidelines for the prevention of cardiovascular disease in women - 2011 update: a guideline from the american heart association. *Circulation.* 2011;123(11):1243-62.
 43. Bechsgaard DF, Hove JD, Suhrs HE, Bové KB, Shahriari P, Gustafsson I, et al. Women with coronary microvascular dysfunction and no obstructive coronary artery disease have reduced exercise capacity. *Int J Cardiol.* 2019;293:1-9.
 44. Suda A, Takahashi J, Beltrame JF, Berry C, Camici PG, Crea F, et al. International prospective cohort study of microvascular angina: rationale and design. *Int J Cardiol Heart Vasc* (in press).
 45. Pristipino C, Beltrame JF, Finocchiaro ML, Hattori R, Fujita M, Mongiardo R, et al. Major racial differences in coronary constrictor response between Japanese and Caucasians with recent myocardial infarction. *Circulation.* 2000;101(10):1102-8.
 46. Ong P, Athanasiadis A, Hill S, Vogelsberg H, Voehringer M, Sechtem U. Coronary artery spasm as a frequent cause of acute coronary syndrome: the CASPAR (Coronary Artery Spasm in Patients With Acute Coronary Syndrome) study. *J Am Coll Cardiol.* 2008;52(7):523-7.
 47. Sato K, Takahashi J, Odaka Y, Suda A, Sueda S, Teragawa H, et al. Clinical characteristics and long-term prognosis of contemporary patients with vasospastic angina: ethnic differences detected in an international comparative study. *Int J Cardiol.* 2019;291:13-8.
 48. JCS Joint Working Group. Guidelines for diagnosis and treatment of patients with hypertrophic cardiomyopathy (JCS 2012). *Circ J.* 2016;80(3):753-74.
 49. Ong P, Athanasiadis A, Borgulya G, Mahrholdt H, Kaski JC, Sechtem U. High prevalence of a pathological response to acetylcholine testing in patients with stable angina pectoris and unobstructed coronary arteries. The ACOVA study (Abnormal COronary VAsomotion in patients with stable angina and unobstructed coronary arteries). *J Am Coll Cardiol.* 2012;59(7):655-62.
 50. Ong P, Pirozzolo G, Athanasiadis A, Sechtem U. Epicardial coronary spasm in women with angina pectoris and unobstructed coronary arteries is linked with a positive family history: an observational study. *Clin Ther.* 201;40(9):1584-90.

51. Kawano H, Motoyama T, Ohgushi M, Kugiyama K, Ogawa H, Yasue H. Menstrual cyclic variation of myocardial ischemia in premenopausal women with variant angina. *Ann Intern Med.* 2001;135(11):977-81.
52. Kawana A, Takahashi J, Takagi Y, Yasuda S, Sakata Y, Tsunoda R, et al. Gender differences in the clinical characteristics and outcomes of patients with vasospastic angina: a report from the Japanese Coronary Spasm Association. *Circ J.* 2013;77(5):1267-74.
53. Takagi Y, Takahashi J, Yasuda S, Miyata S, Tsunoda R, Ogata Y, et al. Prognostic stratification of patients with vasospastic angina: a comprehensive clinical risk score developed by the Japanese Coronary Spasm Association. *J Am Coll Cardiol.* 2013;62(13):1144-53.
54. Kikuchi Y, Yasuda S, Aizawa K, Tsuburaya R, Ito Y, Takeda M, et al. Enhanced Rho-kinase activity in circulating neutrophils of patients with vasospastic angina: a possible biomarker for diagnosis and disease activity assessment. *J Am Coll Cardiol.* 2011;58(12):1231-7.
55. Nihei T, Takahashi J, Hao K, Kikuchi Y, Odaka Y, Tsuburaya R, et al. Prognostic impacts of Rho-kinase activity in circulating leucocytes in patients with vasospastic angina. *Eur Heart J.* 2018;39(11):952-9.
56. Suda A, Takahashi J, Kikuchi Y, Shindo T, Matsumoto Y, Sakata Y, et al. Sex differences in coronary microvascular dysfunction in patients with vasospastic angina. *Circulation.* 2017;136:A17652.
57. Curry RC Jr, Pepine CJ, Sabom MB, Conti CR. Similarities of ergonovine-induced and spontaneous attacks of variant angina. *Circulation.* 1979;59(2):307-12.
58. Sueda S, Miyoshi T, Sasaki Y, Sakaue T, Habara H, Kohno H, et al. Gender differences in sensitivity of acetylcholine and ergonovine to coronary spasm provocation test. *Heart Vessels.* 2016;31(3):322-9.
59. Sato K, Kaikita K, Nakayama N, Horio E, Yoshimura H, Ono T, et al. Coronary vasomotor response to intracoronary acetylcholine injection, clinical features, and long-term prognosis in 873 consecutive patients with coronary spasm: analysis of a single-center study over 20 years. *J Am Heart Assoc.* 2013;2(4):e000227.
60. Takagi Y, Yasuda S, Takahashi J, Tsunoda R, Ogata Y, Seki A, et al. Clinical implications of provocation tests for coronary artery spasm: safety, arrhythmic complications, and prognostic impact: multicentre registry study of the Japanese Coronary Spasm Association. *Eur Heart J.* 2013;34(4):258-67.
61. Nishimiya K, Matsumoto Y, Wang H, Piao Z, Ohyama K, Uzuka H, et al. Absence of adventitial vasa vasorum formation at the coronary segment with myocardial bridge. An optical coherence tomography study. *Int J Cardiol.* 2018; 250:275-7.
62. Aydar Y, Yazici HU, Birdane A, Nasifov M, Nadir A, Ulus T, et al. Gender differences in the types and frequency of coronary artery anomalies. *Tohoku J Exp Med.* 2011;225(4):239-47.
63. Yasue H, Takizawa A, Nagao M, Nishida S, Horie M, Kubota J, et al. Long-term prognosis for patients with variant angina and influential factors. *Circulation.* 1988;78(1): 1-9.
64. Fukumoto Y, Yasuda S, Ito A, Shimokawa H. Prognostic effects of benidipine in patients with vasospastic angina: comparison with diltiazem and amlodipine. *J Cardiovasc Pharmacol.* 2008;51(3):253-7.
65. Beltrame JF, Crea F, Kaski JC, Ogawa H, Ong P, Sechtem U, et al. The Who, What, Why, When, How and Where of Vasospastic Angina. *Circ J.* 2016;80(2):289-98.
66. Takahashi J, Nihei T, Takagi Y, Miyata S, Odaka Y, Tsunoda R, et al. Prognostic impact of chronic nitrate therapy in patients with vasospastic angina: multicentre registry study of the Japanese coronary spasm association. *Eur Heart J.* 2015 Jan 21;36(4):228-37.
67. Ford TJ, Stanley B, Sidik N, Good R, Rocchiccioli P, McEntegart M, et al. 1-year outcomes of Angina Management Guided by Invasive Coronary Function Testing (CorMicA). *J Am Coll Cardiol Interv.* 2020;13(1):33-45.
68. Hiroki J, Shimokawa H, Mukai Y, Ichiki T, Takeshita A. Divergent effects of estrogen and nicotine on Rho-kinase expression in human coronary vascular smooth muscle cells. *Biochem Biophys Res Commun.* 2005;326(1):154-9.
69. Kandabashi T, Shimokawa H, Miyata K, Kunihiro I, Eto Y, Morishige K, et al. Evidence for protein kinase C-mediated activation of Rho-kinase in a porcine model of coronary artery spasm. *Arterioscler Thromb Vasc Biol.* 2003;23(12): 2209-14.
70. Jia H, Abtahian F, Aguirre AD, Lee S, Chia S, Lowe H, et al. In vivo diagnosis of plaque erosion and calcified nodule in patients with acute coronary syndrome by intravascular optical coherence tomography. *J Am Coll Cardiol.* 2013; 62(19):1748-58.
71. Bentzon JF, Otsuka F, Virmani R, Falk E. Mechanisms of plaque formation and rupture. *Circ Res.* 2014;114(12): 1852-66.
72. Farb A, Burke AP, Tang AL, Liang TY, Mannan P, Smialek J, et al. Coronary plaque erosion without rupture into a lipid core. A frequent cause of coronary thrombosis in sudden coronary death. *Circulation.* 1996;93(7):1354-63.

Acknowledgments

We wish to thank the members of the Ischemic Heart Disease Group of the Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, and the staff of the Tohoku University Hospital Cardiac Catheterization Laboratory for their collaboration to the present study.

Funding: this study was supported in part by the grants in aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Author contribution statement: both Authors contributed equally in conceiving, write and review the final version of the manuscript.

Conflict of interest statement: the Authors declare no conflicts of interest.

Correspondence to:

Hiroaki Shimokawa

Department of Cardiovascular Medicine
Tohoku University Graduate School of Medicine
1-1 Seiryomachi, Aoba, Sendai
980-8574 Japan
email shimo@cardio.med.tohoku.ac.jp